

The geochemical behaviour of trace metals along the Belgian coasts: what can we learn from the use of DGT?

Gaulier, Camille; Superville, Pierre-Jean; Guo, Wei; Baeyens, Willy; Billon, Gabriel; Gao, Yue

Published in:
BOOK OF ABSTRACTS - Goldschmidt 2018

Publication date:
2018

Document Version:
Final published version

[Link to publication](#)

Citation for published version (APA):
Gaulier, C., Superville, P.-J., Guo, W., Baeyens, W., Billon, G., & Gao, Y. (2018). The geochemical behaviour of trace metals along the Belgian coasts: what can we learn from the use of DGT? In BOOK OF ABSTRACTS - Goldschmidt 2018: Goldschmidt Abstracts (2018 ed., pp. 799-799). Boston, USA: White Iron Conferences.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

The geochemical behaviour of trace metals along the Belgian coasts: what can we learn from the use of DGT?

CAMILLE GAULIER^{1,2*}, PIERRE-JEAN SUPERVILLE², WEI GUO¹, WILLY BAEYENS¹, GABRIEL BILLON² AND YUE GAO¹

¹Vrije Universiteit Brussel, 1050 Brussels, Belgium

(*correspondence : camille.gaulier@vub.be)

²Lille 1 University, 59655 Villeneuve d'Ascq Cedex, France

Most trace metals exhibit a dual role in marine waters, they act as nutrients in low concentrations, but have toxic effects in the higher concentration ranges. Continuous monitoring efforts in the Belgian Coastal Zone (BCZ) has led to a better understanding of the fate of trace metals in the marine environment. However, the toxicity of these trace elements is strongly linked with their biogeochemical speciation, which shows both seasonal and spatial variation. Furthermore, classical speciation measurements of dissolved vs. particulate trace metals do not correlate well with the bioavailability of these species.

Here, we compare the classical trace metal sampling procedure with an assessment of their bioavailability using Diffusive Gradient in Thin Films (DGT). This dual approach allows to shed light on the relation between trace metal concentrations and the actual bio-available fraction. Based on stable carbon and nitrogen isotope measurements of suspended particulate matter (SPM) we attempt to trace back the origin and identify the anthropogenic fingerprint on the trace metal geochemistry. The BCZ and the Scheldt Estuary are ideal places to study trace metal behaviour, as the proximity of strong industrial activity has led to high concentrations of trace metals, as well as a strong anthropogenic impact. For this study, 13 stations were selected and sampled during several sampling campaigns: 7 stations in the estuary, 4 stations in harbour zones and 2 stations in the North Sea (~ 5km offshore). Our results show that: (i) particulate and total dissolved metal concentrations are higher at estuarine and harbour stations than the offshore ones; (ii) The higher dissolved and particulate concentrations do not correlate with their bioavailability; and (iii) SPM in the harbour zone is likely from allochthonous sources, while in the offshore stations it is mainly from marine origin.

These results indicate that, even though contamination is higher in the harbour zones, the effect of trace metal toxicity (i.e. its bioavailability) is most likely not higher than in the open sea. However, with increasing acidification of the ocean, the higher particulate trace metal concentrations might lead to increasing adverse effects on the coastal environment.