

Editorial: A multi-talented butterfly: the role of the cerebellum in social cognition, emotion, and language

Ma, Qianying; Baetens, Kris; Crunelle, Cleo; Heleven, Elien

Published in:
Frontiers in Human Neuroscience

DOI:
[10.3389/fnhum.2023.1322977](https://doi.org/10.3389/fnhum.2023.1322977)

Publication date:
2023

License:
CC BY

Document Version:
Final published version

[Link to publication](#)

Citation for published version (APA):
Ma, Q., Baetens, K., Crunelle, C., & Heleven, E. (2023). Editorial: A multi-talented butterfly: the role of the cerebellum in social cognition, emotion, and language. *Frontiers in Human Neuroscience*, 17, [1322977]. <https://doi.org/10.3389/fnhum.2023.1322977>

Copyright

No part of this publication may be reproduced or transmitted in any form, without the prior written permission of the author(s) or other rights holders to whom publication rights have been transferred, unless permitted by a license attached to the publication (a Creative Commons license or other), or unless exceptions to copyright law apply.

Take down policy

If you believe that this document infringes your copyright or other rights, please contact openaccess@vub.be, with details of the nature of the infringement. We will investigate the claim and if justified, we will take the appropriate steps.



OPEN ACCESS

EDITED AND REVIEWED BY
Leonhard Schilbach,
Ludwig Maximilian University of
Munich, Germany

*CORRESPONDENCE
Qianying Ma
✉ qianying.ma@vub.be

RECEIVED 17 October 2023
ACCEPTED 18 October 2023
PUBLISHED 03 November 2023

CITATION
Ma Q, Baetens K, Crunelle CL and Heleven E
(2023) Editorial: A multi-talented butterfly: the
role of the cerebellum in social cognition,
emotion, and language.
Front. Hum. Neurosci. 17:1322977.
doi: 10.3389/fnhum.2023.1322977

COPYRIGHT
© 2023 Ma, Baetens, Crunelle and Heleven.
This is an open-access article distributed under
the terms of the [Creative Commons Attribution
License \(CC BY\)](#). The use, distribution or
reproduction in other forums is permitted,
provided the original author(s) and the
copyright owner(s) are credited and that the
original publication in this journal is cited, in
accordance with accepted academic practice.
No use, distribution or reproduction is
permitted which does not comply with these
terms.

Editorial: A multi-talented butterfly: the role of the cerebellum in social cognition, emotion, and language

Qianying Ma^{1,2*}, Kris Baetens², Cleo L. Crunelle³ and
Elien Heleven²

¹Language Pathology and Brain Science MEG Lab, School of Communication Sciences, Beijing Language and Culture University, Beijing, China, ²Brain, Body and Cognition, Center for Neuroscience, Vrije Universiteit Brussel, Brussel, Belgium, ³Department of Psychiatry, Vrije Universiteit Brussel, University Hospital Brussels (UZ Brussel), Brussel, Belgium

KEYWORDS

cerebellum, social cognition, language, emotions, connectivity, psychiatry

Editorial on the Research Topic


A multi-talented butterfly: the role of the cerebellum in social cognition, emotion, and language

In our daily life, understanding what others think, inferring their emotions and communicating with each other can facilitate smooth social interactions. Over the last decades, researchers have been devoted to exploring how the cerebrum processes information in these social interactions. The human cerebellum has a surface that is about 80% of that of the cerebral cortex and has about four times as many neurons (Lent et al., 2012; Sereno et al., 2020). With this in mind, it is surprising that neuroscientists have only relatively recently become interested in cerebellar involvement in social, emotional and linguistic processes (See meta-analyses: Buckner et al., 2011; Van Overwalle et al., 2020a,b, 2023).

Going beyond the traditional understanding of the cerebellum as a motor controller (Manto et al., 2012), there is a growing awareness that the cerebellum also plays a critical role in non-motor domains; from basic cognitive functions to complex social functions. Multiple meta-analyses conducted during the last decade have shown that the cerebellum, especially the posterior cerebellum, is robustly and consistently activated during non-motor tasks (Van Overwalle et al., 2014, 2023; Guell et al., 2018; King et al., 2019).

A recent large-scale meta-analysis examined over 1000 fMRI studies encompassing more than 44,500 participants regarding task-based functional organization within the cerebellum (Van Overwalle et al., 2023). This parcellation suggested that there are 10 functional clusters associated to seven networks, similar to what was proposed by Buckner et al. (2011). As shown in Figure 1, the task-based parcellation demonstrated the involvement of the cerebellum, especially of the posterior cerebellar Crus I and II and the inferior posterior cerebellar IX, in social mentalizing, language and affective task contexts.

Social processes are quite complex, and we often use language and emotions to understand, make inferences and communicate in social spaces with other social beings, which might make it difficult to accurately disentangle these processes in practice. Therefore, the aim of the topic is to bring together findings on the cerebellum's role in social, affective,



and language processing, which can provide a more general idea about how the cerebellum contributes to higher-level processing.

Coemans et al. explored the effects of anodal cerebellar transcranial direct current stimulation (ctDCS) coupled with language therapy in a bilingual patient with chronic post-stroke aphasia caused by left frontal ischemia. They found significant improvements in untrained repetition and picture description tasks in two languages after anodal ctDCS. Additionally, anodal ctDCS led to improved performance on a cognitive control task, suggesting domain-general enhancements in monitoring abilities. Although these are preliminary results, the study opens promising avenues for further research on ctDCS in aphasia and the role of the cerebellum in language processing.

The study by Niu et al. aimed to investigate the neural mechanisms underlying chronic smoking by analyzing spontaneous brain activity and functional connectivity in smokers. The results showed weakened functional connectivity between the prefrontal cortex and cerebellar subregions. This disrupted connectivity may represent a key role of the cerebellum in emotional regulation, attention, and decision-making associated with chronic smoking and addiction.

Dadomo et al. used multivariate pattern analysis to explore the structural neural correlates of borderline personality disorder, a complex mental disorder characterized by unstable emotions, impulsiveness, feelings of inadequacy, and interpersonal problems. The results showed that morphometric differences in the caudate, putamen and amygdala were predictive of sexual trauma, specifically in this population. The results also showed an association between alterations in cerebellar areas and interpersonal problems in patients with borderline personality disorders.

Finally, the study by Jia et al. investigated the neural basis of schizophrenia by examining functional connectivity in the brain using magnetoencephalography imaging. Reduced social skills and positive symptoms (i.e., hallucinations, delusions or repetitive movements) are key symptoms of schizophrenia. The findings showed that, compared to healthy participants, patients with schizophrenia had distorted functional connectivity across delta-theta, alpha, and beta frequency bands. Specifically, there was increased connectivity in beta frequencies between the left primary auditory cortex and cerebellum, linked to more severe hallucinations in schizophrenia. These findings were interpreted in line with the idea that aberrant agency experience in schizophrenia

is the consequence of dysfunctional predictive mechanisms, in turn more broadly in keeping with the view that cerebellar predictive processes are crucially involved in diverse forms of psychopathology (Van Overwalle et al., 2021).

Overall, these studies showed the multifaceted involvement of the cerebellum in a number of high-level processes. Considering the growing consensus that the cerebellum plays a critical role in non-motor domains, it is important to further investigate its contributions. Also, our collected articles showed distorted connectivity between the cerebellum and the corresponding cerebrum in various pathologies, especially those featuring social impairments. Although there is an emerging consensus that cerebellar dysfunction relates to psychiatric disorders, further research is needed to understand this relationship. As a potential target area for brain stimulation, understanding cerebellar functions and its intricate internal and cerebral connections can potentially pave the way for innovative treatments and interventions for clinical conditions related to social processing, emotions and language.

Author contributions

QM: Writing—original draft. KB: Writing—review & editing. CC: Writing—review & editing. EH: Writing—review & editing.

References

- Buckner, R. L., Krienen, F. M., Castellanos, A., Diaz, J. C., and Yeo, B. T. T. (2011). The organization of the human cerebellum estimated by intrinsic functional connectivity. *J. Neurophysiol.* 106, 2322–2345. doi: 10.1152/jn.00339.2011
- Guell, X., Gabrieli, J. D. E., and Schmahmann, J. D. (2018). (2018). Triple representation of language, working memory, social and emotion processing in the cerebellum: convergent evidence from task and seed-based resting-state fMRI analyses in a single large cohort. *NeuroImage* 172, 437–449. doi: 10.1016/j.neuroimage.2018.01.082
- King, M., Hernandez-Castillo, C. R., Poldrack, R. A., Ivry, R. B., and Diedrichsen, J. (2019). Functional boundaries in the human cerebellum revealed by a multi-domain task battery. *Nat. Neurosci.* 22, 1371–1378. doi: 10.1038/s41593-019-0436-x
- Lent, R., Azevedo, F. A. C., Andrade-Moraes, C. H., and Pinto, A. V. O. (2012). How many neurons do you have? Some dogmas of quantitative neuroscience under revision. *Eur. J. Neurosci.* 35, 1–9. doi: 10.1111/j.1460-9568.2011.07923.x
- Manto, M., Bower, J. M., Conforto, A. B., Delgado-García, J. M., Guarda, D., A., et al. (2012). Consensus paper: roles of the cerebellum in motor control—the diversity of ideas on cerebellar involvement in movement. *Cerebellum* 11, 457–487. doi: 10.1007/s12311-011-0331-9
- Sereno, M. I., Diedrichsen, J., Tachrount, M., Testa-Silva, G. D., Arceuil, H., and De Zeeuw, C. (2020). The human cerebellum has almost 80% of the surface area of the neocortex. *Proc. Nat. Acad. Sci. U. S. A.* 117, 19538–19543. doi: 10.1073/pnas.2002896117
- Van Overwalle, F., Baeken, C., Campanella, S., Crunelle, C. L., Heleven, E., Kornreich, C., et al. (2021). The role of the posterior cerebellum in dysfunctional social sequencing. *The Cereb.* 21, 1–12. doi: 10.1007/s12311-021-01330-y
- Van Overwalle, F., Baetens, K., Mariën, P., and Vandekerckhove, M. (2014). Social cognition and the cerebellum: a meta-analysis of over 350 fMRI studies. *NeuroImage* 86, 554–572. doi: 10.1016/j.neuroimage.2013.09.033
- Van Overwalle, F., Ma, Q., Haihambo, N., Bylemans, T., Catoira, B., Firouzi, M., et al. (2023). A functional atlas of the cerebellum based on neurosynth task coordinates. *Cerebellum*. 22, 1–20. doi: 10.1007/s12311-023-01596-4
- Van Overwalle, F., Ma, Q., and Heleven, E. (2020a). The posterior crus II cerebellum is specialized for social mentalizing and emotional self-experiences: a meta-analysis. *Soc. Cognit. Aff. Neurosci.* 15, 905–928. doi: 10.1093/scan/nsaa124
- Van Overwalle, F., Manto, M., Cattaneo, Z., Clausi, S., Ferrari, C., Gabrieli, J. D. E., et al. (2020b). Consensus paper: cerebellum and social cognition. *Cerebellum* 19, 833–868. doi: 10.1007/s12311-020-01155-1

Funding

This research project was supported by Science Foundation of Beijing Language and Culture University (supported by the Fundamental Research Funds for the Central Universities) (23YJ110001) awarded to QM. Also, this research project was supported by Strategic Research Program SRP57 from the Vrije Universiteit Brussel.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.