

Interim safety and efficacy results of a phase II clinical trial on trametinib and low-dose dabrafenib in patients with advanced BRAFV600 wild-type melanoma

Awada, Gil; Schwarze, Julia Katharina; Reijmen, Eva; Goyvaerts, Cleo; Fasolino, Giuseppe; Aspeslagh, Sandrine; Neyns, Bart

Published in:
Annals of Oncology

Publication date:
2020

Document Version:
Final published version

[Link to publication](#)

Citation for published version (APA):

Awada, G., Schwarze, J. K., Reijmen, E., Goyvaerts, C., Fasolino, G., Aspeslagh, S., & Neyns, B. (2020). Interim safety and efficacy results of a phase II clinical trial on trametinib and low-dose dabrafenib in patients with advanced BRAFV600 wild-type melanoma. *Annals of Oncology*, 31(S1), S10-S10.

Copyright

No part of this publication may be reproduced or transmitted in any form, without the prior written permission of the author(s) or other rights holders to whom publication rights have been transferred, unless permitted by a license attached to the publication (a Creative Commons license or other), or unless exceptions to copyright law apply.

Take down policy

If you believe that this document infringes your copyright or other rights, please contact openaccess@vub.be, with details of the nature of the infringement. We will investigate the claim and if justified, we will take the appropriate steps.

190 A phase Ib trial of CFI-402257 in combination with weekly paclitaxel in patients with advanced HER2-negative (HER2-) breast cancer (aBC)

P. Bedard¹, M. Mates², K.A. Gelmon³, J. Hilton⁴, D. Tu⁵, I. Li¹, L. Rastgou⁶, N. Drummond-Ivars⁴, J. Edwards⁵, D. Warr¹, C. Lohrisch⁷, X. Song⁴, A. Srikanthan⁴, A. Robinson², L. Hagerman⁵, L.K. Seymour⁸, M. Rushton⁸, D.W. Cescon¹

¹Princess Margaret Cancer Center, Toronto, ON, Canada; ²Cancer Centre of South-eastern Ontario, Kingston, ON, Canada; ³Medical Oncology Department, BC Cancer, Vancouver, BC, Canada; ⁴Medical Oncology Dept, The Ottawa Hospital Regional Cancer Centre, Ottawa, ON, Canada; ⁵Canadian Clinical Trials Group, Kingston, ON, Canada; ⁶Medical Oncology Department, BC Cancer Agency - Vancouver, Vancouver, BC, Canada; ⁷BC Cancer Agency - Vancouver, Vancouver, BC, Canada; ⁸IND Program, Canadian Cancer Trials Group, Kingston, ON, Canada

Background: CFI-402257 is a selective oral inhibitor of TTK protein kinase, a critical regulator of the mitotic spindle assembly checkpoint. TTK is overexpressed in breast cancer (BC); CFI-402257 has anti-proliferative and cytotoxic activity as monotherapy. Synergy has been identified between TTK inhibitors and taxanes; CFI-402257 enhances the antitumor activity of paclitaxel in BC xenograft models.

Methods: The primary objective was to establish the safety and recommended phase 2 dose (RP2D) of CFI-402257 combined with weekly paclitaxel. Patients with HER2-ve aBC with adequate organ function, PS 0-1, previously treated with > 1 non-taxane chemotherapy, not appropriate for endocrine therapy, were eligible. A 3+3 design was used with dose limiting toxicities (DLTs) assessed during cycle 1 (28 days). Starting dose CFI-402257 was 84mg on 2-day on, 5-day off schedule with paclitaxel 80mg/m² day 1, 8, and 15. Safety assessments were performed weekly (CTCAE v5.0) and response (RECIST 1.1) every 2 cycles.

Results: 13 patients received a total of 39 cycles at 3 dose levels (84mg, 112mg, 168mg). Median age was 51, 85% ER+/HER2-ve, 54% PS1, 50% ≥3 prior lines of chemotherapy, and 54% with ≥4 sites of metastatic disease. One grade 4 neutropenia DLT was seen at DL3; no related serious adverse events (AEs) were seen. The most frequent (≥30%) non-hematological AEs were alopecia (82%), diarrhea (55%), nausea (55%), fatigue (55%), vomiting (45%), sensory neuropathy (45%), headache (45%), constipation (45%), back pain (36%), and extremity pain (36%). Grade ≥3 hematological AEs were neutropenia (27%) and anemia (9%). The overall response rate for evaluable patients was 1/9 = 11.1%.

Conclusions: CFI-402257 and paclitaxel is well tolerated. Dose escalation is ongoing and the RP2D has not yet been defined. Updated safety and efficacy data will be presented at the meeting.

Clinical trial identification: NCT03568422.

Legal entity responsible for the study: Canadian Cancer Trials Group.

Funding: This study was coordinated by the Canadian Cancer Trials Group supported by SU2C Canada - Canadian Cancer Society Breast Cancer Dream Team Research Funding (SU2C-AAACR-DT-18-15), with supplemental support from the Ontario Institute for Cancer Research, through funding provided by the Government of Ontario.

Disclosure: P. Bedard: Research grant/Funding (institution): AstraZeneca; Research grant/Funding (institution): Novartis; Advisory/Consultancy, Research grant/Funding (institution): BMS; Research grant/Funding (institution): GSK; Research grant/Funding (institution): Seattle Genetics; Research grant/Funding (institution): Merck; Advisory/Consultancy, Research grant/Funding (institution): Sanofi; Research grant/Funding (institution): Pfizer; Research grant/Funding (institution): Lilly; Research grant/Funding (institution): Servier; Research grant/Funding (institution): PTC Therapeutics; Research grant/Funding (institution): SignalChem LifeSciences; Research grant/Funding (institution): Genentech/Roche; Research grant/Funding (institution): Nektar Therapeutics; Research grant/Funding (institution): Zymeworks. K.A. Gelmon: Advisory/Consultancy, Research grant/Funding (institution): AZ; Advisory/Consultancy, Research grant/Funding (institution): Roche; Advisory/Consultancy, Research grant/Funding (institution): Novartis; Advisory/Consultancy: Pfizer; Advisory/Consultancy: Oncothyreon; Advisory/Consultancy: Nanostring; Advisory/Consultancy: Merck; Advisory/Consultancy: Mylan; Advisory/Consultancy: Eli Lilly; Advisory/Consultancy: Genomic Health; Advisory/Consultancy, Research grant/Funding (institution): BMS; Speaker Bureau/Expert testimony: Genentech. J. Hilton: Advisory/Consultancy: Pfizer; Advisory/Consultancy: Lilly; Advisory/Consultancy: Puma; Advisory/Consultancy: Novartis; Advisory/Consultancy, Non-remunerated activity/ies, Data Monitoring Committee: BMS; Advisory/Consultancy: AZ. L.K. Seymour: Research grant/Funding (institution), For trial costs: University Health Network. D.W. Cescon: Advisory/Consultancy, Research grant/Funding (institution): Merck; Advisory/Consultancy, Research grant/Funding (institution): Pfizer; Advisory/Consultancy, Research grant/Funding (institution): GSK; Advisory/Consultancy: Novartis; Advisory/Consultancy: AZ; Advisory/Consultancy: Roche; Advisory/Consultancy: Puma; Advisory/Consultancy: Agendia; Advisory/Consultancy: Dynamo Therapeutics; Non-remunerated activity/ies, Patent holder (to institution) for biomarkers related to CFI-402257: University Health Network. All other authors have declared no conflicts of interest.

<https://doi.org/10.1016/j.annonc.2020.01.028>

20P Interim safety and efficacy results of a phase II clinical trial on trametinib and low-dose dabrafenib in patients with advanced BRAFV600 wild-type melanoma

G. Awada¹, J.K. Schwarze¹, E. Reijmen², C. Goyvaerts², G. Fasolino³, S. Aspeslagh¹, B. Neyns¹

¹Department of Medical Oncology, Universitair Ziekenhuis Brussel, Brussels, Belgium; ²Laboratory of Molecular & Cellular Therapy, Vrije Universiteit Brussel, Brussels, Belgium; ³Department of Ophthalmology, Universitair Ziekenhuis Brussel, Brussels, Belgium

Background: Preclinical and clinical data suggest MEK-inhibition (MEKi) to be effective in NRAS-mutant (mt) and NRAS wild-type (wt) melanoma. However, MEKi causes considerable cutaneous treatment-related adverse events (TRAE) which are less present if MEKi is combined with BRAFi.

Methods: This open-label, dual-stratum, single-center phase 2 clinical trial investigated trametinib (T; 2 mg QD) in patients (pts) with advanced BRAF^{V600} wt, NRAS^{C61R/K/L} mt/wt melanoma who had progressed following or were ineligible for immune checkpoint inhibitors. In case of T-related cutaneous TRAE, low-dose dabrafenib (Id-D; 50 mg BID) was added to prevent further skin TRAE. The trial was amended in June 2019 to administer Id-D upfront with T. The primary endpoint was the confirmed objective response rate (ORR; per RECIST 1.1); secondary endpoints were progression-free and overall survival and safety.

Results: Between Jan and Dec 2019, 9 pts initiated T monotherapy and 8 pts initiated T + Id-D. TRAE were seen in 15 pts (29% G3-4; 29% SAE). One pt permanently interrupted T due to G3 pneumonitis. All 9 pts who initiated T monotherapy developed G1-2 acneiform rash which resolved (to G0: 6 pts; to G1: 3 pts) with temporary interruption of T, local metronidazole therapy and subsequent addition of Id-D to T. One pt had a G1 recurrence of acneiform rash that was managed with local therapy. Of 8 pts who initiated T + Id-D upfront, 1 developed G1 acneiform rash that resolved with local therapy only. Other TRAE are shown in the table. T dose was reduced in 4 pts for TRAE (1 pt with left ventricular ejection fraction decrease; 1 with central serous retinopathy; 1 with AST/ALT increase; 1 with hyponatremia and syncope). The unconfirmed ORR is 4/15 and the disease control rate is 7/15 evaluable pts.

Table 20P: TRAE in ≥2 pts

TRAE	G1-2 (n[%])	G3-4 (n[%])
CPK increase	10 (59)	0
Diarrhea	5 (29)	0
AST increase	4 (24)	1 (6)
Lipase increase	4 (24)	0
Arterial hypertension	4 (24)	0
Hyponatremia	2 (12)	2 (12)
ALT increase	3 (18)	1 (6)
Fatigue	2 (12)	1 (6)
Anemia	3 (18)	0
Chills	3 (18)	0
Nausea	3 (18)	0
Central serous retinopathy	3 (18)	0
Syncope	0	2 (12)
Acute kidney injury	2 (12)	0
Thrombocytopenia	2 (12)	0
Fever	2 (12)	0

Conclusions: No unexpected toxicities were seen with T or T + Id-D. Id-D is able to prevent T-related skin toxicity. Thus, combining Id-D with T is a safe approach to increase tolerance of and optimize exposure to T.

Clinical trial identification: NCT04059224; EudraCT 2018-004003-39.

Legal entity responsible for the study: Universitair Ziekenhuis Brussel.

Funding: Novartis, Stichting tegen Kanker.

Disclosure: G. Awada: Research grant/Funding (institution), Travel/Accommodation/Expenses: Pfizer; Travel/Accommodation/Expenses: Astellas; Travel/Accommodation/Expenses: Merck Sharp & Dohme; Advisory/Consultancy, Research grant/Funding (institution): Novartis. J.K. Schwarze: Travel/Accommodation/Expenses: Amgen; Travel/Accommodation/Expenses: Merck Sharp & Dohme. S. Aspeslagh: Advisory/Consultancy, Speaker Bureau/Expert testimony: Amgen; Advisory/Consultancy, Speaker Bureau/Expert testimony: Roche; Advisory/Consultancy, Speaker Bureau/Expert testimony: AstraZeneca; Advisory/Consultancy, Speaker Bureau/Expert testimony: Bristol-Myers Squibb; Advisory/Consultancy, Speaker Bureau/Expert testimony: Merck Sharp & Dohme; Advisory/Consultancy, Speaker Bureau/Expert testimony: Sanofi; Advisory/Consultancy, Speaker Bureau/Expert testimony: Novartis. B. Neyns: Advisory/Consultancy, Speaker Bureau/Expert testimony: Amgen; Advisory/Consultancy, Speaker Bureau/Expert testimony: Bristol-Myers Squibb; Advisory/Consultancy, Speaker Bureau/Expert testimony: Novartis; Advisory/Consultancy, Speaker Bureau/Expert testimony: Merck Sharp & Dohme; Advisory/Consultancy, Speaker Bureau/Expert testimony: AstraZeneca; Advisory/Consultancy, Speaker Bureau/Expert testimony: EthernA; Advisory/Consultancy, Speaker Bureau/Expert testimony: CryoStorage; Research grant/Funding (institution): Pfizer; Research grant/Funding (institution): Merck Serono. All other authors have declared no conflicts of interest.

<https://doi.org/10.1016/j.annonc.2020.01.039>