

High CO₂ content in magmas at an arc volcano: the andesitic Enco eruption of Mocho-Choshuenco volcano (Chile)

Feignon, Jean-Guillaume; Roche, Olivier; Moune, Séverine; Cluzel, Nicolas; Schiavi, Federica; Clavero, Jorge; Schiano, Pierre; Auxerre, Marion

Published in:
EGU General Assembly Conference Abstracts 2019

Publication date:
2019

License:
CC BY

Document Version:
Final published version

[Link to publication](#)

Citation for published version (APA):
Feignon, J-G., Roche, O., Moune, S., Cluzel, N., Schiavi, F., Clavero, J., Schiano, P., & Auxerre, M. (2019). High CO₂ content in magmas at an arc volcano: the andesitic Enco eruption of Mocho-Choshuenco volcano (Chile). In *EGU General Assembly Conference Abstracts 2019: EGU2019-16065* (Vol. 21, pp. 16065-16065). Geophysical Research Abstracts.

Copyright

No part of this publication may be reproduced or transmitted in any form, without the prior written permission of the author(s) or other rights holders to whom publication rights have been transferred, unless permitted by a license attached to the publication (a Creative Commons license or other), or unless exceptions to copyright law apply.

Take down policy

If you believe that this document infringes your copyright or other rights, please contact openaccess@vub.be, with details of the nature of the infringement. We will investigate the claim and if justified, we will take the appropriate steps.



High CO₂ content in magmas at an arc volcano: the andesitic Enco eruption of Mocho-Choshuenco volcano (Chile)

Jean-Guillaume Feignon (1,2), Olivier Roche (1), Séverine Moune (1), Nicolas Cluzel (1), Federica Schiavi (1), Jorge Clavero (3), Pierre Schiano (1), and Marion Auxerre (4)

(1) Laboratoire Magmas et Volcans, Université Clermont Auvergne, CNRS, IRD, OPGC, F-63000 Clermont-Ferrand, France (jean-guillaume.feignon@etudiant.univ-bpclermont.fr), (2) Department of Lithospheric Research, University of Vienna, Althanstraße 14, A-1090 Vienna, Austria, (3) Escuela de Geología, Universidad Mayor, Santiago, Chile, (4) Centre de Recherches Péetrographiques et Géochimiques (CRPG), UMR CNRS 5873, Université de Lorraine, BP20, 54501 Vandœuvre les Nancy, France

High volatile contents are thought to be one of the main causes of explosivity of poorly to moderately differentiated magmas that lead to effusive volcanic eruptions otherwise. In particular, CO₂ may be a key factor in promoting high gas pressures and subsequent magma fragmentation owing to its low solubility.

We address this issue by investigating the pre-eruptive volatile (H₂O, CO₂, Cl and S) content in the crystal-poor andesitic magma that led to the sub-Plinian Enco eruption of Mocho-Choshuenco volcano, one of the most hazardous volcanoes in the southern volcanic zone of Chile.

Though the whole rock composition of the Enco magma is andesitic (60.2±1.1 wt.% SiO₂), the melt inclusions have compositions ranging from 50.3 to 67.3 wt.% SiO₂, following the magmatic series of Mocho-Choshuenco, and the compositions of the most mafic melt inclusions are close to that of the most mafic erupted magmas.

We measured the volatile contents in melt inclusions trapped in minerals (olivine, plagioclase and pyroxene) using electron microprobe and ion microprobe (SIMS). Glass analysis revealed typical parental arc magma values for H₂O (2.6-3.1 wt.%), S (116-1936 ppm) and Cl (620-1993 ppm), but exceptionally high contents in CO₂ with concentrations above 4000 ppm in some melt inclusions. Presence of solid carbonates inside inclusion-hosted bubbles, revealed by 3D confocal Raman mapping, clearly indicated that the CO₂ contents measured in the glass phase were minimum values. We conclude that the high CO₂ content could be one of the causes of explosivity of the Enco eruption that generated pyroclastic flows.